

兔抗 STMN1(Phospho-Ser62) 多克隆抗体

中文名称：兔抗 STMN1(Phospho-Ser62) 多克隆抗体

英文名称： Anti-STMN1(Phospho-Ser62) rabbit polyclonal antibody

别名： Lag; SMN; OP18; PP17; PP19; PR22; LAP18; C1orf215

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： STMN1(Phospho-Ser62)

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

| | |
|--------------------------|---|
| Background: | This gene belongs to the stathmin family of genes. It encodes a ubiquitous cytosolic phosphoprotein proposed to function as an intracellular relay integrating regulatory signals of the cellular environment. The encoded protein is involved in the regulation of the microtubule filament system by destabilizing microtubules. It prevents assembly and promotes disassembly of microtubules. Alternatively spliced transcript variants encoding the same protein have been identified. |
| Applications: | WB, IHC |
| Name of antibody: | STMN1(Phospho-Ser62) |

| | |
|---------------------------------|---|
| Immunogen: | Peptide sequence around phosphorylation site of Serine 62(R-R-K(p)-S-H) derived from Human STMN1. |
| Full name: | stathmin 1 |
| Synonyms : | Lag; SMN; OP18; PP17; PP19; PR22; LAP18; C1orf215 |
| SwissProt: | P16949 |
| IHC positive control: | Human breast carcinoma tissue |
| IHC Recommend dilution: | 50-100 |
| WB Predicted band size: | 17 kDa |
| WB Positive control: | COS cells lysates treated with nocodazole |
| WB Recommended dilution: | 500-1000 |



